

Minimal Textures in Seesaw Mass Matrices and their low and high Energy Phenomenology

Srubabati Goswami^{a,b*}, Subrata Khan^{a†}, Werner Rodejohann^{c‡}

^a*Physical Research Laboratory,
Navrangpura, Ahmedabad 380009, India*

^b*Harish-Chandra Research Institute,
Chhatnag Road, Jhansi, Allahabad 211019, India*

^c*Max-Planck-Institut für Kernphysik,
Postfach 103980, D-69029 Heidelberg, Germany*

Abstract

In an attempt to find minimal scenarios we study the implications of Dirac and Majorana mass matrices with texture zeros within the type I seesaw mechanism. For the Dirac mass matrices we consider 5 zero textures which we show to be the most minimal form that can successfully account for low energy phenomenology if the Majorana mass matrices are chosen minimal as well. For those, we consider both diagonal and even more minimal non-diagonal forms. The latter can be motivated e.g. by simple $U(1)$ flavour symmetries and have two degenerate eigenvalues. We classify the allowed textures and discuss the ramifications for leptogenesis and lepton flavour violation.

*email: sruba@prl.res.in

†email: subrata@prl.res.in

‡email: werner.rodejohann@mpi-hd.mpg.de

1 Introduction

Spectacular results from neutrino oscillation experiments in the past decade have established the existence of neutrino masses and lepton mixing on firm footing. However, the smallness of the neutrino masses still continues to be an enigma. The most elegant mechanism for generating small neutrino masses is the seesaw mechanism in which one adds heavy right-handed singlets (type I) [1], scalar triplets (type II) [2] or fermion triplets (type III) [3] to generate small neutrino masses at low scale. In the context of the type I seesaw mechanism the mass matrix for the left handed neutrinos obtained through seesaw diagonalization depends on the Dirac type Yukawa coupling matrix of the neutrinos (m_D) as well as on the bare Majorana mass matrix (M_R) of the heavy right handed neutrinos:

$$m_\nu = -m_D M_R^{-1} m_D^T. \quad (1.1)$$

Reconstruction of seesaw mass matrices from low energy observations is a challenging task. The main problem is the mismatch in the number of parameters because in general the seesaw framework contains more parameters compared to what can be obtained from measurements at low energy and it is not possible to fix the high energy parameters entirely from low energy data [4] (for an overview, see [5]). One possible solution is the appearance of "texture zeros". In general, "zeros" imply vanishingly small entries in the mass matrices whose origin can be traced to flavour symmetries. Consideration of texture zeros in the seesaw mass matrices provides an useful way to handle the problem of parameter mismatch since assumption of texture zeros leads to a reduction of the number of parameters at high scale and thus strengthens the predictive power of the model.

Implications of two texture zeros in the low energy Majorana mass matrix have been studied in [6, 7, 8] and one texture zero have been studied in [9] (see also [10] for the case of additional equality of mass matrix entries). Texture zeros in both the charged lepton and neutrino mass matrices have been studied in [11].

Within the framework of seesaw mechanism it is often considered more natural to consider texture zeros appearing in the Yukawa coupling matrix m_D and/or the right-handed Majorana mass matrix M_R [12, 13, 14]. In particular, it has been shown in [13] that if M_R is diagonal and if one assumes that all light neutrino states are massive then the maximal number of zeros that can be accommodated in m_D is four. The phenomenology of those cases is studied in detail in Ref. [14]. We will relax this assumption here since the low energy data allow one of the neutrino states to be massless. In that case, as we show, even with a diagonal M_R one can obtain allowed textures consistent with low energy observations for a m_D with 5 zeros. The results in [13] were obtained assuming both the charged lepton mass matrix and the heavy neutrino mass matrix to be diagonal. We also relax the assumption of a diagonal M_R . However, we insist in non-singular M_R in order to avoid light right-handed (sterile) neutrinos. With this constraint we find that the maximal number of zeros that is possible for M_R is four and only three such matrices have a non-vanishing determinant. Interestingly these matrices obey $L_\mu - L_\tau$, $L_e - L_\tau$ and $L_e - L_\mu$ flavour symmetry. We investigate if it is possible to have allowed textures assuming the above forms for M_R and m_D with 5 zeros, and find that such textures are disallowed by low

energy constraints. It is easy to convince oneself that more zeros in m_D are not allowed for the minimal forms of M_R that we study [15]. To complete the survey we consider the implications for Lepton Flavour Violation (LFV) and leptogenesis for the allowed textures surviving the constraints from low energy phenomenology.

The plan of the paper goes as follows: in the next Section we discuss the seesaw phenomenology including its manifestation for leptogenesis and LFV. In Section 3 we perform the texture analysis for m_D with 5 zeros for both diagonal M_R (3 zeros) and non-diagonal M_R with 4 zeros and obeying simple flavour symmetries. The results in terms of allowed and disallowed textures are presented in Section 4. Apart from low energy phenomenology, we also discuss the implications for leptogenesis and LFV for the allowed solutions. Finally we make some comments on Dirac matrices with more than 5 zeros and conclude in Section 5.

2 Seesaw Phenomenology

The low energy neutrino mass matrix given in Eq. (1.1) is symmetric and can in general be diagonalized with a unitary matrix as

$$V_\nu^T m_\nu V_\nu = D_\nu, \quad (2.1)$$

where D_ν is diagonal, real and positive and contains the mass eigenvalues for the left-handed neutrinos: $D_\nu = \text{diag}(m_1, m_2, m_3)$. In the basis in which the charged lepton mass matrix is real and diagonal, V_ν coincides with the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix U and the standard parametrization for this matrix is:

$$U = \begin{pmatrix} c_{12} c_{13} & s_{12} c_{13} & s_{13} e^{-i\delta} \\ -c_{23} s_{12} - s_{23} s_{13} c_{12} e^{i\delta} & c_{23} c_{12} - s_{23} s_{13} s_{12} e^{i\delta} & s_{23} c_{13} \\ s_{23} s_{12} - c_{23} s_{13} c_{12} e^{i\delta} & -s_{23} c_{12} - c_{23} s_{13} s_{12} e^{i\delta} & c_{23} c_{13} \end{pmatrix} P. \quad (2.2)$$

Here $c_{ij} = \cos \theta_{ij}$, $s_{ij} = \sin \theta_{ij}$, δ is the Dirac-type CP-violating phase and the Majorana phases α and β are contained in the matrix $P = \text{diag}(1, e^{i\alpha}, e^{i(\beta+\delta)})$. While all phases are currently unconstrained, the other mixing parameters are determined with increasing precision. Table 1 summarizes the results from Ref. [16].

Neutrino oscillation experiments are sensitive to the Dirac CP-phase δ . Instead of the phase δ it is often more common to use the rephasing invariant quantity J_{CP} (Jarlskog invariant)

$$J_{CP} = \text{Im} \{ U_{e1} U_{\mu 2} U_{e2}^* U_{\mu 1}^* \} = -\frac{\text{Im} \{ h_{12} h_{23} h_{31} \}}{\Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2}, \text{ where } h = m_\nu m_\nu^\dagger. \quad (2.3)$$

With the parameterization of Eq. (2.2), one has $J_{CP} = \frac{1}{8} \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \sin \delta$.

	$\Delta m_{\odot}^2/10^{-5} \text{ eV}^2$	$\sin^2 \theta_{12}$	$ U_{e3} $	$\sin^2 \theta_{23}$	$\Delta m_{\text{A}}^2/10^{-3} \text{ eV}^2$
Best-fit	7.67	0.312	0.126	0.466	2.39
1σ	$7.48 \div 7.83$	$0.294 \div 0.331$	$0.077 \div 0.161$	$0.408 \div 0.539$	$2.31 \div 2.50$
3σ	$7.14 \div 8.19$	$0.263 \div 0.375$	< 0.214	$0.331 \div 0.644$	$2.06 \div 2.81$

Table 1: Current best-fit values as well as 1 and 3σ ranges of the oscillation parameters [16].

Another place where CP violation in the lepton sector plays a useful role is leptogenesis, even though in general the low and high energy CP violations can be completely uncorrelated. Both leptogenesis and the later to be discussed Lepton Flavour Violation (LFV) need to be evaluated in the basis in which M_R is real and diagonal with $M_1 \leq M_2 \leq M_3$. In general the Majorana mass matrix M_R is non-diagonal in the basis where the charged current is flavour diagonal. It is thus written as

$$U_R^\dagger M_R U_R^* = \text{diag}(M_1, M_2, M_3). \quad (2.4)$$

After performing a basis rotation so that the right-handed Majorana mass matrix M_R becomes diagonal by the unitary matrix U_R , the Dirac mass matrix m_D gets modified to

$$m_D \rightarrow \tilde{m}_D = m_D U_R^*. \quad (2.5)$$

Considering the decay of one heavy Majorana neutrino N_i into the Higgs and lepton doublets, the CP asymmetry generated through the interference between tree level and one loop heavy Majorana neutrino decay diagrams is given as [17]

$$\begin{aligned} \varepsilon_i^\alpha &\equiv \frac{\Gamma(N_i \rightarrow \phi \bar{l}_\alpha) - \Gamma(N_i \rightarrow \phi^\dagger l_\alpha)}{\sum_\beta \left[\Gamma(N_i \rightarrow \phi \bar{l}_\beta) + \Gamma(N_i \rightarrow \phi^\dagger l_\beta) \right]} \\ &= \frac{1}{8\pi v_u^2} \frac{1}{(\tilde{m}_D^\dagger \tilde{m}_D)_{ii}} \sum_{j \neq i} \left(\mathcal{I}_{ij}^\alpha f(M_j^2/M_i^2) + \mathcal{J}_{ij}^\alpha \frac{1}{1 - M_j^2/M_i^2} \right), \end{aligned} \quad (2.6)$$

where

$$\mathcal{I}_{ij}^\alpha = \text{Im} \left[(\tilde{m}_D^\dagger)_{i\alpha} (\tilde{m}_D)_{\alpha j} (\tilde{m}_D^\dagger \tilde{m}_D)_{ij} \right], \quad \mathcal{J}_{ij}^\alpha = \text{Im} \left[(\tilde{m}_D^\dagger)_{i\alpha} (\tilde{m}_D)_{\alpha j} (\tilde{m}_D^\dagger \tilde{m}_D)_{ji} \right]. \quad (2.7)$$

In the MSSM, the function $f(x)$ has the form [17]

$$f(x) = \sqrt{x} \left[\frac{2}{1-x} - \ln \left(\frac{1+x}{x} \right) \right]. \quad (2.8)$$

Here ε_i^α describes the decay asymmetry of the right-handed neutrino of mass M_i into leptons of flavour $\alpha = e, \mu, \tau$. If the rest-mass of the lightest heavy neutrino is much lighter than

the other two, i.e. $M_1 \ll M_{2,3}$, the lepton asymmetry is dominated by the decay of this lightest of the heavy neutrinos. In this case $f(M_j^2/M_1^2) \simeq -3 M_1/M_j$. Moreover, only the first term, proportional to $\mathcal{I}_{1j}^\alpha$, in Eq. (2.6) is relevant since the second term proportional to $\mathcal{J}_{ij}^\alpha$ is suppressed by an additional power of M_1/M_j . Let us mention here that we will find all $\mathcal{J}_{1j}^\alpha$ to be zero for the successful textures. Note furthermore that the second term in Eq. (2.6) vanishes when one sums over flavours to obtain the flavour independent decay asymmetry:

$$\begin{aligned} \varepsilon_i &= \sum_\alpha \varepsilon_i^\alpha \equiv \frac{\sum_\alpha \left[\Gamma(N_i \rightarrow \phi \bar{l}_\alpha) - \Gamma(N_i \rightarrow \phi^\dagger l_\alpha) \right]}{\sum_\beta \left[\Gamma(N_i \rightarrow \phi \bar{l}_\beta) + \Gamma(N_i \rightarrow \phi^\dagger l_\beta) \right]} \\ &= \frac{1}{8\pi v_u^2} \frac{1}{(\tilde{m}_D^\dagger \tilde{m}_D)_{ii}} \sum_{j \neq i} \text{Im} \left[(\tilde{m}_D^\dagger \tilde{m}_D)_{ij}^2 \right] f(M_j^2/M_i^2) \\ &= \frac{1}{8\pi v_u^2} \frac{1}{(\tilde{m}_D^\dagger \tilde{m}_D)_{ii}} \mathcal{I}_{ij}, \text{ where } \mathcal{I}_{ij} = \sum_\alpha \mathcal{I}_{ij}^\alpha. \end{aligned} \quad (2.9)$$

The expressions given above for the decay asymmetries are valid for the MSSM. Their flavour structure is, however, identical to that of just the Standard Model.

Equally important in leptogenesis are effective mass parameters that are responsible for the wash-out. With our assumption that a single heavy neutrino of mass M_1 is relevant for leptogenesis, the wash-out of every decay asymmetry ε_1^α is governed by an effective mass

$$\tilde{m}_1^\alpha = \frac{(\tilde{m}_D^\dagger)_{1\alpha} (\tilde{m}_D)_{\alpha 1}}{M_1}. \quad (2.10)$$

The summation of $\tilde{m}_1^\alpha$ over the flavour index α yields $\tilde{m}_1$, which is the relevant parameter for the wash-out of ε_1 . One needs to insert the effective masses in the function

$$\eta(x) \simeq \left(\frac{8.25 \times 10^{-3} \text{ eV}}{x} + \left(\frac{x}{2 \times 10^{-4} \text{ eV}} \right)^{1.16} \right)^{-1}. \quad (2.11)$$

The final baryon asymmetry is [18, 19, 20]

$$Y_B \simeq \begin{cases} -0.01 \varepsilon_1 \eta(\tilde{m}_1) & \text{one-flavour,} \\ -0.003 \left((\varepsilon_1^e + \varepsilon_1^\mu) \eta \left(\frac{417}{589} (\tilde{m}_1^e + \tilde{m}_1^\mu) \right) + \varepsilon_1^\tau \eta \left(\frac{390}{589} \tilde{m}_1^\tau \right) \right) & \text{two-flavour,} \\ -0.003 \left(\varepsilon_1^e \eta \left(\frac{151}{179} \tilde{m}_1^e \right) + \varepsilon_1^\mu \eta \left(\frac{344}{537} \tilde{m}_1^\mu \right) + \varepsilon_1^\tau \eta \left(\frac{344}{537} \tilde{m}_1^\tau \right) \right) & \text{three-flavour.} \end{cases} \quad (2.12)$$

Here we have given separate expressions for one-, two- and three-flavoured leptogenesis. The three-flavour case occurs for $M_1 (1 + \tan^2 \beta) \leq 10^9 \text{ GeV}$, the one-flavour case for $M_1 (1 + \tan^2 \beta) \geq 10^{12} \text{ GeV}$, and the two-flavour case (with the tau-flavour decoupling first and the sum of electron- and muon-flavours, which act indistinguishably) applies in between.

Another possible part of seesaw phenomenology is Lepton Flavour Violation (LFV) in supersymmetric seesaw scenarios. Decays such as $\ell_i \rightarrow \ell_j \gamma$, with flavour indices i, j spanning ($1 = e, 2 = \mu, 3 = \tau$) can depend on the same parameters, though in a different form, as the light neutrino masses. In mSUGRA scenarios with universal boundary conditions for scalar sparticle mass matrices, renormalization effects from M_X down to the scale of the heavy neutrino masses induce off-diagonal entries in the slepton mass matrix [21]:

$$\text{BR}(\ell_i \rightarrow \ell_j + \gamma) = c \times \text{BR}(\ell_i \rightarrow \ell_j \nu \bar{\nu}) |(\tilde{m}_D L \tilde{m}_D^\dagger)_{ij}|^2, \quad (2.13)$$

where the diagonal matrix L is defined as

$$L_{kl} = \ln \frac{M_X}{M_k} \delta_{kl}, \quad (2.14)$$

with M_k being the mass of the k^{th} right-handed neutrino. The constant c in the RHS of Eq. (2.13) depends on certain supersymmetry parameters of mSUGRA, specifically the universal scalar and gaugino masses and the universal trilinear scalar coupling as well as on $\tan \beta$. However, we are not interested here in the exact magnitude of the branching ratios. We shall instead study the vanishing of certain branching ratios or the ratio of branching ratios, which is independent on the constant.

3 Texture Analysis

3.1 Criteria for successful Candidates and their Phenomenology

Once we have evaluated the low energy neutrino mass matrix from m_D and M_R containing zero entries, it turns out that invalid cases can easily be ruled almost at first sight.

First of all, the rank of m_ν should be at least two. The resulting m_ν will have zeros in most of the cases. A very powerful result obtained in [6] is that low energy Majorana mass matrices with more than 2 zeros are not consistent with data*. From the 15 possible two zero textures† only seven are found to be consistent with the data. The allowed patterns are characterized by the simultaneous vanishing of the ee - and $e\mu$ -, the ee - and $e\tau$ -, the $e\mu$ - and $\mu\mu$ -, the $e\tau$ - and $\mu\mu$ -, the $e\tau$ - and $\tau\tau$ -, the $e\mu$ - and $\tau\tau$ -, and finally the $\mu\mu$ - and $\tau\tau$ -entries. The first two possibilities are only possible for a normal hierarchy, the latter five only for quasi-degenerate neutrinos. In particular, if e.g. one neutrino mass is zero then it is not allowed that $(m_\nu)_{\mu\mu} = (m_\nu)_{\tau\tau} = 0$. If the $\mu\tau$ entry of m_ν is zero, then no other entry is allowed to vanish. Regarding the presence of only one zero entry in m_ν , an important information is that if the determinant of m_ν vanishes, then the $\mu\tau$ element can not be zero [9].

*Let us note here that seesaw realizations of low energy two zero textures have been analyzed in Ref. [7].

†If neutrino are Dirac particles then up to five zero entries are allowed in m_ν [22].

Another necessary result to take into account is the “scaling” behavior of m_ν [23, 24, 25]. This denotes the following form of the mass matrix:

$$m_\nu = \begin{pmatrix} a & b & b/c \\ \cdot & d & d/c \\ \cdot & \cdot & d/c^2 \end{pmatrix}. \quad (3.1)$$

This form necessarily occurs *regardless of the form of M_R* if the third row of m_D multiplied with c is equal to the second row [25]. In the context of texture zeros, this means that each of the second and third row of m_D contain only one entry in the same column. Alternatively, the second row is zero and the third row contains only one non-zero element. The low energy matrix in Eq. (3.1) is only compatible with an inverted hierarchy, $m_3 = U_{e3} = 0$ and $\tan^2 \theta_{23} = 1/c^2$. Solar neutrino mixing is unspecified by the above matrix, and the effective mass governing neutrino-less double beta decay is $\langle m \rangle = \sqrt{\Delta m_A^2} \sqrt{1 - \sin^2 2\theta_{12} \sin^2 \alpha}$. There are in principle other variants of scaling, which will result in m_ν similar to Eq. (3.1), but with the first and second, or first and third column correlated. These are incompatible with data and can be identified already by the form of m_D .

The simple set of properties summarized here is basically all what is needed to distinguish the allowed possibilities from the disallowed ones. A somewhat more tricky situation is dealt with in Section 4.2.

Let us also comment on the possible impact of the Renormalization Group (RG) effects [26] on the allowed textures. The matrix m_ν that we obtain through seesaw diagonalization is at a high scale determined by the masses of the heavy neutrinos. Therefore it is expected that the low scale predictions will be affected in general by the RG effects. However the impact is in general small for hierarchical mass spectrum[‡]. It is also well known that the RG effects on the mass matrix m_ν are multiplicative in nature. Therefore we expect a zero to remain a zero even after RG evolution. Thus, an allowed texture is expected to be stable against RG corrections [28]. One can also show that mass matrices with the scaling property are stable against RG effects [23].

Finally, we would like to mention that we do not take into account the fine-tuned possibility that the low energy texture zeros are resulting from cancellation of terms.

3.2 M_R more minimal than diagonal

Heavy neutrino mass matrices in diagonal form contain three independent zeros. One can become even more minimal by choosing the following, non-singular matrices with four

[‡]Considerable running even for normal hierarchy can be obtained if running in between the mass scales of the individual heavy neutrinos is taken into account [27]. In order to have our conclusions and findings stable against such effects, we therefore need to assume that the zeros are protected by the same symmetry which is responsible for them.

independent zeros:

$$M_R^{-1} = \begin{pmatrix} M_1 & 0 & 0 \\ 0 & 0 & M_2 \\ 0 & M_2 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 & M_2 \\ 0 & M_1 & 0 \\ M_2 & 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & M_2 & 0 \\ M_2 & 0 & 0 \\ 0 & 0 & M_1 \end{pmatrix}. \quad (3.2)$$

Interestingly, these correspond to flavour symmetries $L_\mu - L_\tau$, $L_e - L_\tau$ and $L_e - L_\mu$, respectively [§]. Majorana (and thus symmetric) matrices with 5 or more zeros are all singular, lead to light sterile neutrinos and will not be studied here. As we will see, an exactly diagonal M_R leads to allowed forms of m_ν either with scaling or with single zero entries. Using the above textures of M_R will give typically two simultaneously vanishing entries in m_ν .

With m_D being a in general non-symmetric 3×3 matrix, there are $N = {}^9C_n$ possibilities to place n zero entries in it. We therefore have e.g. $N = 126, 84$ and 36 if $n = 5, 6$ and 7 . With our analysis aiming at 5 zero textures in m_D and with the four forms of M_R , there are in total 504 candidates. Most of them can be ruled out by the arguments given in Section 3.1.

4 Results

In what follows we classify the allowed textures of m_D and M_R in terms of the allowed forms of the low energy Majorana mass matrix m_ν .

4.1 Diagonal M_R

If M_R is diagonal then we find in total 18 allowed cases, all of which generate a vanishing determinant for m_ν :

- (i) six of them have a vanishing $e\mu$ entry in m_ν . The cases are listed in Table 2.
- (ii) another six contain a vanishing $e\tau$ element. The cases are listed in Table 3.
- (iii) the remaining six fulfill the scaling condition in Eq. (3.1). The cases are listed in Table 4.

Let us first discuss the neutrino mixing phenomenology. The predictions of mass matrices obeying scaling were given after Eq. (3.1). The other 12 cases are specified by a vanishing element in m_ν and $m_1 = 0$ or $m_3 = 0$, depending on whether a normal or inverted hierarchy is present. In case of $(m_\nu)_{e\mu} = 0$ it has been shown that in general θ_{13} must necessarily be non-zero [9]. If $m_1 = 0$ then one finds from the condition $(m_\nu)_{e\mu} = 0$ that

$$|U_{e3}| \simeq \frac{m_2 \cos \theta_{12} \cot \theta_{23}}{\sqrt{\Delta m_A^2 / \sin^2 \theta_{12} - 2\sqrt{\Delta m_A^2 \Delta m_\odot^2} \cos 2\alpha}} \simeq \begin{cases} 0.077 \div 0.113 & \text{at } 1\sigma \\ 0.050 \div 0.144 & \text{at } 3\sigma \end{cases}. \quad (4.1)$$

[§]Note that the M_R corresponding to the matrices in 3.2 will be obtained by just replacing M_i by $1/M_i$. Thus in this convention the M_i s in equation 3.2 and throughout the text will have inverse mass dimension.

The numerical range of $|U_{e3}|$ is obtained by using the exact expression for $(m_\nu)_{e\mu}$ and varying the neutrino parameters in their currently allowed 1 and 3σ ranges. In an inverted hierarchy, with $m_3 = 0$ and $m_2 \simeq m_1$ we find

$$|U_{e3}| \simeq \frac{\sin 2\theta_{12} \sin \alpha}{\sqrt{1 - \sin^2 2\theta_{12} \sin^2 \alpha}} \cot \theta_{23}. \quad (4.2)$$

While in principle $|U_{e3}|$ can be sizable according to this equation, it turns out [9] that the vanishing of the $e\mu$ -element implies also that $|\sin \alpha| \ll 1$. In fact, expanding in terms of $|U_{e3}|$ one finds up to first order that

$$(m_\nu)_{e\mu} = (m_2 e^{2i\alpha} - m_1) \cos \theta_{12} \sin \theta_{12} \cos \theta_{23} + e^{i\delta} \sin \theta_{23} (e^{2i\beta} m_3 - \sin^2 \theta_{12} m_2 e^{2i\alpha} - \cos^2 \theta_{12} m_1) |U_{e3}|, \quad (4.3)$$

We see that for $m_2 \simeq m_1 \gg m_3$ the phase should be such that $e^{2i\alpha} \simeq 1$. This value implies for the effective mass that $\langle m \rangle \simeq \sqrt{\Delta m_A^2} \cos^2 \theta_{13}$, i.e., there are basically no cancellations. We find numerically a lower value of $|U_{e3}| \gtrsim 0.005$ when the 3σ ranges of the oscillation parameters are used, whereas for the 1σ ranges we get $|U_{e3}| \gtrsim 0.082$. If the $e\tau$ -entry of m_ν is zero, then similar expressions as Eqs. (4.1) and (4.2) are obtained, the only change to be made is that $\cot \theta_{23}$ should be replaced with $\tan \theta_{23}$, which has little numerical consequences.

Turning to leptogenesis, we see from Tables 2, 3 and 4 that for all 18 cases there is only one physical phase present. In addition, there is always only one flavoured $\mathcal{I}_{1j}^\alpha$ present (which is equal to the unflavoured $\mathcal{I}_{1j}$) and the corresponding wash-out parameter $\tilde{m}_1^\alpha$ (which is equal to the unflavoured $\tilde{m}_1$) is non-zero. Note that in some cases m_D is such that the neutrino with mass M_1 decouples. Then we have calculated the corresponding quantities for the second neutrino, i.e. $\mathcal{I}_{2j}^\alpha$ and $\tilde{m}_2^\alpha$. As indicated, all $\mathcal{J}_{1j}^\alpha$ are zero. For all 18 possibilities there is the potential to accommodate successful leptogenesis.

We have checked that for the cases with a vanishing $e\mu$ or $e\tau$ element in m_ν the invariant J_{CP} from Eq. (2.3), responsible for low energy CP violation in oscillations is non-zero and is proportional to the same phase factor as in $\mathcal{I}_{ij}$. For instance, if we consider

$$m_D = \begin{pmatrix} 0 & 0 & a_3 \\ 0 & b_2 & 0 \\ 0 & c_2 & c_3 e^{i\gamma_3} \end{pmatrix}, \quad (4.4)$$

which gives m_ν with $m_{e\mu} = 0$ for a diagonal M_R , then the only non-zero $\mathcal{I}_{ij}$ that we get is

$$\mathcal{I}_{23} = c_2^2 c_3^2 \sin 2\gamma_3, \quad (4.5)$$

to be compared with

$$J_{\text{CP}} = a_3^2 b_2^2 c_2^2 c_3^2 (a_3^2 (b_2^2 + c_2^2) + b_2^2 c_3^2) M_2^3 M_3^3 \sin 2\gamma_3. \quad (4.6)$$

Thus we see that $\mathcal{I}_{ij}^\tau$ and J_{CP} contain the same phase factor $\sin 2\gamma_3$. Hence, the Dirac CP phase is always identical to the high-energy “leptogenesis phase”. This conclusion is true for all the textures in Tables 2 and 3.

An exception is when scaling in m_ν is present, where there is no CP violation in oscillation experiments. Here it may be helpful to evaluate the invariant [29]

$$I_{e\mu} = \text{Im} \left\{ (m_\nu)_{ee} (m_\nu)_{\mu\mu} (m_\nu)_{e\mu}^* (m_\nu)_{\mu e}^* \right\} .$$

In general, this expression is rather lengthy, but we can express it in simple form for the case of $m_3 = 0$ and $U_{e3} = 0$. Inserting these constraints in the standard parametrization of U from Eq. (2.2) and evaluating the invariant yields $I_{e\mu} = -m_1 m_2 \Delta m_\odot^2 c_{12}^2 s_{12}^2 c_{23}^4 \sin 2\alpha$ in terms of the low energy parameters. Similarly $I_{e\tau}$ is also non-zero and depends on the same phase. $I_{\mu\tau}$ is however zero. Note that α corresponds to the Majorana phase, the only physical low energy phase if scaling is present. In terms of the parameters of m_D and M_R , $I_{e\mu}$ can be expressed in terms of the single phase that characterizes the Dirac mass matrices. For instance if we consider

$$m_D = \begin{pmatrix} 0 & a_2 & a_3 e^{i\alpha_3} \\ 0 & 0 & b_3 \\ 0 & 0 & c_3 \end{pmatrix} \quad (4.7)$$

from Table 4, which gives m_ν obeying the scaling property, then

$$I_{e\mu} = a_2^2 a_3^2 b_3^4 M_2 M_3^3 \sin 2\alpha_3 . \quad (4.8)$$

We note from Table 4 that the same phase factor $\sin 2\alpha_3$ from $I_{e\mu}$ also appears in $\mathcal{I}_{23}^e$. Hence, for this case we can say that the Majorana CP phase crucial for neutrino-less double beta decay is identical to the high-energy “leptogenesis phase”. This conclusion is true for all the textures in Table 4. Finally, it is worth remarking that the fifth entry in Table 4 corresponds to the results of a model based on the flavour symmetry $D_4 \times Z_2$ constructed in Ref. [23].

Let us note that in the case of 4 zero textures [13, 14] there always was an ambiguity in what regards the one-to-one identification of low and high energy CP phases with each other. This is not the case for 5 zero textures.

Regarding LFV, in case the $e\mu$ entry of m_ν vanishes, then also $(m_D L m_D^\dagger)_{12}$ is zero. Hence, $\mu \rightarrow e\gamma$ will not be detected if the only source of LFV is the supersymmetric seesaw with mSUGRA conditions studied here. The remaining, in general non-zero branching ratios turn out to be proportional to the respective elements of m_ν . Consider the first matrix in Table 2. The low energy mass matrix is

$$m_\nu = - \begin{pmatrix} \frac{a_3^2}{M_3} & 0 & \frac{a_3 c_3 e^{i\gamma_3}}{M_3} \\ \cdot & \frac{b_2^2}{M_2} & \frac{b_2 c_2}{M_2} \\ \cdot & \cdot & \frac{c_2^2}{M_2} + \frac{c_3^2 e^{i\gamma_3}}{M_3} \end{pmatrix} , \quad (4.9)$$

and for LFV we find

$$|(m_D L m_D^\dagger)_{13}|^2 = a_3^2 c_3^2 L_3^2, \quad |(m_D L m_D^\dagger)_{23}|^2 = b_2^2 c_2^2 L_2^2. \quad (4.10)$$

Hence, $\text{BR}(\tau \rightarrow e\gamma) \propto |(m_\nu)_{e\tau}|^2$ and $\text{BR}(\tau \rightarrow \mu\gamma) \propto |(m_\nu)_{\mu\tau}|^2$. The ratio of $\text{BR}(\tau \rightarrow e\gamma)$ and $\text{BR}(\tau \rightarrow \mu\gamma)$ can however not be predicted in general because the a priori unknown heavy neutrino masses enter this ratio directly and via the factors L_i . The qualitatively same situation is encountered for a vanishing $e\tau$ entry of m_ν , in which case $\text{BR}(\tau \rightarrow e\gamma)$ is zero and the other BR are proportional to the respective elements of m_ν , but with different dependence on the heavy masses.

In contrast, if m_ν obeys scaling, then no off-diagonal element of $(m_D L m_D^\dagger)_{ij}$ vanishes. Consider the first matrix in Table 4, for which

$$m_\nu = - \begin{pmatrix} \frac{a_2^2}{M_2} + \frac{a_3^2 e^{2i\alpha_3}}{M_3} & \frac{a_3 b_3 e^{i\alpha_3}}{M_3} & \frac{a_3 c_3 e^{i\alpha_3}}{M_3} \\ \cdot & \frac{b_3^2}{M_3} & \frac{b_3 c_3}{M_3} \\ \cdot & \cdot & \frac{c_3^2}{M_3} \end{pmatrix} \quad (4.11)$$

and

$$|(m_D L m_D^\dagger)_{12}|^2 = a_3^2 b_3^2 L_3^2, \quad |(m_D L m_D^\dagger)_{13}|^2 = a_3^2 c_3^2 L_3^2, \quad |(m_D L m_D^\dagger)_{23}|^2 = b_3^2 c_3^2 L_3^2.$$

In particular, we can make the definite prediction

$$\frac{|(m_D L m_D^\dagger)_{12}|^2}{|(m_D L m_D^\dagger)_{13}|^2} = \frac{b_3^2}{c_3^2} = \cot^2 \theta_{23}. \quad (4.12)$$

This relation is in fact true in every seesaw model leading to scaling [23]. A property of the textures is that the branching ratios are proportional to their respective mass matrix entries. We can calculate

$$\frac{|(m_\nu)_{e\mu}|^2}{|(m_\nu)_{\mu\tau}|^2} = \frac{\text{BR}(\mu \rightarrow e\gamma) \text{BR}(\tau \rightarrow \mu \nu \bar{\nu})}{\text{BR}(\tau \rightarrow \mu\gamma)} \simeq \frac{\sin^2 2\theta_{12} \sin^2 \alpha}{\sin^2 \theta_{23} (1 - \sin^2 2\theta_{12} \sin^2 \alpha)}. \quad (4.13)$$

Here, we have neglected $\Delta m_\odot^2$ with respect to Δm_A^2 . This ratio is naturally of order one, but can also vanish. Numerically, for the 1σ (3σ) ranges of the oscillation parameter the ratio is less than roughly 18.2 (42.9). If $\alpha = 0$, then it becomes $(\Delta m_\odot^2 / \Delta m_A^2)^2 \sin^2 2\theta_{12} / (16 \sin^2 \theta_{23})$. Note that Eq. (4.12) implies that $\tau \rightarrow e\gamma$ will be too rare to become observable.

Finally, it is worth mentioning if a μ - τ symmetric seesaw is possible. This would imply

$$m_D = \begin{pmatrix} a_1 & a_2 & a_2 \\ b_1 & b_2 & b_3 \\ b_1 & b_2 & b_3 \end{pmatrix}, \quad M_R = \begin{pmatrix} M_1 & 0 & 0 \\ 0 & M_2 & 0 \\ 0 & 0 & M_2 \end{pmatrix}. \quad (4.14)$$

This situation would lead to a μ - τ symmetric low energy mass matrix of the form

$$m_\nu = \begin{pmatrix} a & b & b \\ \cdot & d & f \\ \cdot & \cdot & d \end{pmatrix} \quad (4.15)$$

with $U_{e3} = 0$ and $\sin^2 \theta_{23} = \frac{1}{2}$. It can be seen from Tables 2, 3 and 4 that none of the allowed 5 zero textures is compatible with μ - τ symmetry. At low energy we can disregard this possibility at once for the cases of a zero $(m_\nu)_{e\mu}$ or a zero $(m_\nu)_{e\tau}$ entry. Namely, it would imply that also $(m_\nu)_{e\tau}$ and $(m_\nu)_{e\mu}$ are zero, which is not allowed by data. Regarding the matrices obeying scaling, there can be “accidental” μ - τ symmetry at low energy, namely if $\tan^2 \theta_{23} = 1$. However, this is not a consequence of μ - τ symmetry of the seesaw mass matrices m_D and M_R .

4.2 Non-diagonal M_R

If M_R takes on the non-diagonal four zero textures in Eq. (3.2) then we can generate two zero textures in m_ν . However, only four of the seven allowed two zero textures can be obtained. For each of the three possibilities $L_e - L_\mu$, $L_e - L_\tau$ and $L_\mu - L_\tau$, at first sight 8 potentially successful cases survive:

- for every non-diagonal form of M_R there are two cases with a vanishing ee and $e\mu$ entry in m_ν . They have the additional property

$$|(m_\nu)_{\mu\mu} (m_\nu)_{\tau\tau}| = |(m_\nu)_{\mu\tau}^2| \quad \text{and} \quad \arg \left\{ (m_\nu)_{\mu\mu} (m_\nu)_{\tau\tau} ((m_\nu)_{\mu\tau}^*)^2 \right\} = 0; \quad (4.16)$$

- for every non-diagonal form of M_R there are two cases with a vanishing ee and $e\tau$ entry in m_ν . They have the same property given in Eq. (4.16);
- for every non-diagonal form of M_R there are two cases with a vanishing $e\mu$ and $\mu\mu$ entry in m_ν . They have the additional property

$$|(m_\nu)_{ee} (m_\nu)_{\tau\tau}| = |(m_\nu)_{e\tau}^2| \quad \text{and} \quad \arg \left\{ (m_\nu)_{ee} (m_\nu)_{\tau\tau} ((m_\nu)_{e\tau}^*)^2 \right\} = 0; \quad (4.17)$$

- for every non-diagonal form of M_R there are two cases with a vanishing $e\tau$ and $\tau\tau$ entry in m_ν . They are subject to the condition

$$|(m_\nu)_{ee} (m_\nu)_{\mu\mu}| = |(m_\nu)_{e\mu}^2| \quad \text{and} \quad \arg \left\{ (m_\nu)_{ee} (m_\nu)_{\mu\mu} ((m_\nu)_{e\mu}^*)^2 \right\} = 0; \quad (4.18)$$

All 24 m_ν have a non-vanishing determinant, hence no neutrino mass is zero. They also share the property of a zero invariant J_{CP} , i.e., in the parametrization of the PMNS matrix applied here the CP phase δ is zero or π . However, it is known [6, 8] that if $(m_\nu)_{e\mu} = (m_\nu)_{\mu\mu} = 0$ or if $(m_\nu)_{e\tau} = (m_\nu)_{\tau\tau} = 0$, the CP phase should be large because the small

ratio of solar and atmospheric Δm^2 is proportional to $\cos \delta$. Hence, these 12 cases can be disregarded at once.

In the cases with $(m_\nu)_{ee} = (m_\nu)_{e\mu} = 0$ and $(m_\nu)_{ee} = (m_\nu)_{e\tau} = 0$, it is extremely cumbersome to try to obtain analytical estimates from these equations. However, we have checked that no point in the allowed parameter space can simultaneously satisfy the two zero texture conditions and conditions (4.17) and (4.18). The reason for this lies in a clash of the two zero textures which require hierarchical neutrino masses, whereas the additional conditions imply larger neutrino masses.

4.3 Dirac Mass Matrices with more than five Zeros

Finally, in this Subsection we shortly discuss Dirac mass matrices with more than five zero entries.

Among the 9 elements of m_D which are in general complex the first non-trivial case with maximal number of zeros is 8. This implies 2 heavy neutrinos are completely decoupled from the light neutrinos and it is well known that one cannot explain the low energy phenomenology successfully with only one heavy neutrino coupled to the system.

If m_D contains seven zeros, then we find no valid m_ν for the four minimal forms of M_R under study. In fact we have checked that the above statement is true for the most general form of M_R . One can group the ${}^7C_9 = 36$ possible forms of m_D , in 3 categories. Their respective structures are

$$m_D = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & X \\ 0 & 0 & X \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & X & X \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & X & 0 \\ 0 & 0 & X \end{pmatrix}. \quad (4.19)$$

The first two matrices will lead, irrespective of M_R , in the seesaw mechanism to at least two vanishing eigenvalues in m_ν and are thus ruled out in general. The third one can give two non-vanishing eigenvalues but we have checked that the final m_ν contains more than 2 zeros for the most general form of M_R and hence is disallowed.

For the case of 6 zeros in m_D the possible types of patterns are

$$\begin{pmatrix} 0 & 0 & X \\ 0 & 0 & X \\ 0 & 0 & X \end{pmatrix}, \quad \begin{pmatrix} X & 0 & 0 \\ 0 & X & 0 \\ 0 & 0 & X \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ X & X & X \end{pmatrix}, \quad \begin{pmatrix} X & 0 & 0 \\ 0 & X & 0 \\ X & 0 & 0 \end{pmatrix}. \quad (4.20)$$

For m_D 's of the first and third form, two eigenvalues of m_ν are zero for a general M_R . Hence these forms of m_D are ruled out. For m_D 's of the second type none of the eigenvalues of m_ν are zero and these cannot be ruled out for a general M_R . But we have checked that for the 4 types of M_R that we consider here, the resultant m_ν contains more than 2 zeros and hence are ruled out. For the m_D 's of the fourth type one eigenvalue of m_ν is zero but it contains more than 2 zeros in the flavour basis for a general M_R and therefore are not allowed.

Thus we conclude that for diagonal M_R the 5 zero textures in m_D correspond to scenarios with a minimal number of free parameters. In total, we have a seesaw scenario with 8 zero entries, and if the charged leptons are also taken into account there are in total 14 zeros.

5 Conclusions and Summary

In this paper we have investigated the implications of Dirac mass matrices (m_D) with 5 zeros for seesaw phenomenology. For the Majorana mass matrices (M_R) we consider both diagonal and non-diagonal forms. The diagonal form contains three zeros. However we noted that more minimal forms of M_R containing 4 zeros are possible. Three non-singular forms of M_R exist, which correspond to $L_\mu - L_\tau$, $L_e - L_\tau$ and $L_e - L_\mu$ symmetry. However, with 5 zero textures in m_D they can be shown to be incompatible with neutrino data. We have classified the allowed textures and discussed their implications for leptogenesis and lepton flavour violation. For m_D with 5 zeros and diagonal M_R there are 18 allowed textures

- (i) 6 patterns with a vanishing $e\mu$ entry;
- (ii) 6 patterns with a vanishing $e\tau$ entry;
- (iii) 6 patterns obeying the scaling property with the 2nd and 3rd columns of m_ν proportional to each other.

All these 18 cases have one zero mass eigenvalue. The phenomenology of the cases (i) and (ii) is quite similar: θ_{13} is necessarily non-zero and sizable (at 3σ between 0.05 and 0.14 for the normal hierarchy and larger than 0.005 in the inverted hierarchy) and δ is in general non-trivial. Both normal and inverted hierarchy is possible. Matrices from category (iii) satisfying the scaling property imply $U_{e3} = \delta = 0$ and the hierarchy is inverted. Leptogenesis is possible for all 18 patterns and for the first 12 cases the same phase is responsible for leptogenesis as well as low energy CP violation in neutrino oscillations. For the 6 allowed textures adhering to the scaling property the leptogenesis phase turns out to be the same as the Majorana phase responsible for possible cancellations in neutrinoless double beta decay.

We have also considered the manifestations of seesaw mechanism for LFV by considering supersymmetric seesaw with mSUGRA conditions as the only source for LFV. For the patterns with $(m_\nu)_{e\mu} = 0$ the $\text{BR}(\mu \rightarrow e\gamma)$ turns out to be zero. The $\text{BR}(\tau \rightarrow e\gamma)$ and $\text{BR}(\tau \rightarrow \mu\gamma)$ are proportional to $(m_\nu)_{e\tau}$ and $(m_\nu)_{\mu\tau}$, respectively, but depend on the unknown heavy neutrino masses and hence no definite predictions are possible for these. Similar conclusions hold true for patterns with $(m_\nu)_{e\tau} = 0$. For mass matrices obeying scaling properties definite predictions for LFV in terms of oscillation parameters can be made.

In conclusion, we have shown that with the minimal forms of the Majorana mass matrices that we have chosen (non-diagonal with 4 zeros and diagonal corresponding to 3

zeros) 5 is the maximal number of zeros in Dirac matrices that can generate successful low energy phenomenology.

Acknowledgments

W.R. wishes to thank the Physical Research Laboratory in Ahmedabad and the organizers of the workshop “Astroparticle Physics – A Pathfinder to New Physics” in Stockholm, where part of this paper was written, for hospitality. This work was supported by the ERC under the Starting Grant MANITOP and by the Deutsche Forschungsgemeinschaft in the Transregio 27 (W.R.). S.G. acknowledges partial support from the Neutrino Project under the XI-th plan of Harish-Chandra Research Institute.

References

- [1] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe* (O. Sawada and A. Sugamoto, eds.), KEK, Tsukuba, Japan, 1979, p. 95; M. Gell-Mann, P. Ramond, and R. Slansky, *Complex spinors and unified theories*, in *Supergravity* (P. van Nieuwenhuizen and D. Z. Freedman, eds.), North Holland, Amsterdam, 1979, p. 315; S. L. Glashow, *The future of elementary particle physics*, in *Proceedings of the 1979 Cargèse Summer Institute on Quarks and Leptons* (M. Lévy, J.-L. Basdevant, D. Speiser, J. Weyers, R. Gastmans, and M. Jacob, eds.), Plenum Press, New York, 1980, pp. 687–713. R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [2] M. Magg and C. Wetterich, Phys. Lett. B **94**, 61 (1980); G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); R. N. Mohapatra and G. Senjanovic, Phys. Rev. D **23**, 165 (1981); J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980); J. Schechter and J. W. F. Valle, Phys. Rev. D **25**, 774 (1982).
- [3] R. Foot, H. Lew, X. G. He and G. C. Joshi, Z. Phys. C **44**, 441 (1989).
- [4] S. Davidson and A. Ibarra, JHEP **0109**, 013 (2001) [arXiv:hep-ph/0104076]; J. R. Ellis and M. Raidal, Nucl. Phys. B **643**, 229 (2002) [arXiv:hep-ph/0206174]; S. Pascoli, S. T. Petcov and W. Rodejohann, Phys. Rev. D **68**, 093007 (2003) [arXiv:hep-ph/0302054].
- [5] W. Rodejohann, Pramana **72**, 217 (2009) [arXiv:0804.3925 [hep-ph]].
- [6] P. H. Frampton, S. L. Glashow and D. Marfatia, Phys. Lett. B **536**, 79 (2002) [arXiv:hep-ph/0201008].
- [7] A. Kageyama, S. Kaneko, N. Shimoyama and M. Tanimoto, Phys. Lett. B **538**, 96 (2002).

- [8] Z. Z. Xing, Phys. Lett. B **530**, 159 (2002) [arXiv:hep-ph/0201151]; Z. Z. Xing, Phys. Lett. B **539**, 85 (2002) [arXiv:hep-ph/0205032]; B. R. Desai, D. P. Roy and A. R. Vaucher, Mod. Phys. Lett. A **18**, 1355 (2003) [arXiv:hep-ph/0209035]; C. Hagedorn, J. Kersten and M. Lindner, Phys. Lett. B **597**, 63 (2004) [arXiv:hep-ph/0406103]; S. Dev, S. Kumar, S. Verma and S. Gupta, Phys. Rev. D **76**, 013002 (2007) [arXiv:hep-ph/0612102]; S. Dev, S. Kumar, S. Verma and S. Gupta, Phys. Lett. B **656**, 79 (2007) [arXiv:0708.3321 [hep-ph]]; A. Dighe and N. Sahu, arXiv:0812.0695 [hep-ph].
- [9] A. Merle and W. Rodejohann, Phys. Rev. D **73**, 073012 (2006) [arXiv:hep-ph/0603111].
- [10] S. Kaneko, H. Sawanaka and M. Tanimoto, JHEP **0508**, 073 (2005) [arXiv:hep-ph/0504074].
- [11] Z. Z. Xing and H. Zhang, Phys. Lett. B **569**, 30 (2003) [arXiv:hep-ph/0304234]. Z. Z. Xing, Int. J. Mod. Phys. A **19**, 1 (2004) [arXiv:hep-ph/0307359]. S. Zhou and Z. Z. Xing, Eur. Phys. J. C **38**, 495 (2005) [arXiv:hep-ph/0404188]. Z. Z. Xing and S. Zhou, Phys. Lett. B **593**, 156 (2004) [arXiv:hep-ph/0403261]. M. Randhawa, G. Ahuja and M. Gupta, Phys. Lett. B **643**, 175 (2006) [arXiv:hep-ph/0607074].
- [12] G. K. Leontaris, S. Lola, C. Scheich and J. D. Vergados, Phys. Rev. D **53**, 6381 (1996) [arXiv:hep-ph/9509351]; S. M. Barr and I. Dorsner, Nucl. Phys. B **585**, 79 (2000) [arXiv:hep-ph/0003058]; A. Kageyama, S. Kaneko, N. Shimoyama and M. Tanimoto, Phys. Lett. B **538**, 96 (2002) [arXiv:hep-ph/0204291]; P. H. Frampton, S. L. Glashow and T. Yanagida, Phys. Lett. B **548**, 119 (2002) [arXiv:hep-ph/0208157]; R. Barbieri, T. Hambye and A. Romanino, JHEP **0303**, 017 (2003) [arXiv:hep-ph/0302118]; A. Ibarra and G. G. Ross, Phys. Lett. B **591**, 285 (2004) [arXiv:hep-ph/0312138]; S. Chang, S. K. Kang and K. Siyeon, Phys. Lett. B **597**, 78 (2004) [arXiv:hep-ph/0404187]; A. Watanabe and K. Yoshioka, JHEP **0605**, 044 (2006) [arXiv:hep-ph/0601152]; W. L. Guo, Z. Z. Xing and S. Zhou, Int. J. Mod. Phys. E **16**, 1 (2007) [arXiv:hep-ph/0612033]; S. Goswami and A. Watanabe, Phys. Rev. D **79**, 033004 (2009) [arXiv:0807.3438 [hep-ph]]; S. Goswami, S. Khan and A. Watanabe, arXiv:0811.4744 [hep-ph];
- [13] G. C. Branco, D. Emmanuel-Costa, M. N. Rebelo and P. Roy, Phys. Rev. D **77**, 053011 (2008) [arXiv:0712.0774 [hep-ph]].
- [14] S. Choubey, W. Rodejohann and P. Roy, Nucl. Phys. B **808**, 272 (2009) [arXiv:0807.4289 [hep-ph]].
- [15] See also the reference by A. Watanabe and K. Yoshioka in [12].
- [16] G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, Phys. Rev. Lett. **101**, 141801 (2008) [arXiv:0806.2649 [hep-ph]]; arXiv:0809.2936 [hep-ph].

- [17] L. Covi, E. Roulet and F. Vissani, Phys. Lett. B **384**, 169 (1996).
- [18] A. Abada *et al.*, JCAP **0604**, 004 (2006) [arXiv:hep-ph/0601083];
- [19] E. Nardi, Y. Nir, E. Roulet and J. Racker, JHEP **0601**, 164 (2006) [arXiv:hep-ph/0601084].
- [20] A. Abada, S. Davidson, A. Ibarra, F. X. Josse-Michaux, M. Losada and A. Riotto, JHEP **0609**, 010 (2006) [arXiv:hep-ph/0605281].
- [21] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57**, 961 (1986); J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. D **53**, 2442 (1996) [arXiv:hep-ph/9510309].
- [22] C. Hagedorn and W. Rodejohann, JHEP **0507**, 034 (2005) [arXiv:hep-ph/0503143].
- [23] R. N. Mohapatra and W. Rodejohann, Phys. Lett. B **644**, 59 (2007) [arXiv:hep-ph/0608111]; A. Blum, R. N. Mohapatra and W. Rodejohann, Phys. Rev. D **76**, 053003 (2007) [arXiv:0706.3801 [hep-ph]];
- [24] W. Grimus and L. Lavoura, J. Phys. G **31**, 683 (2005) [arXiv:hep-ph/0410279]; W. Grimus and L. Lavoura, Phys. Rev. D **62**, 093012 (2000) [arXiv:hep-ph/0007011]; L. Lavoura, Phys. Rev. D **62**, 093011 (2000) [arXiv:hep-ph/0005321]; M. S. Berger and S. Santana, Phys. Rev. D **74**, 113007 (2006) [arXiv:hep-ph/0609176]; A. Damanik, M. Satriawan, Muslim and P. Anggraita, arXiv:0705.3290 [hep-ph]; M. Obara, arXiv:0712.2628 [hep-ph];
- [25] A. S. Joshipura and W. Rodejohann, arXiv:0905.2126 [hep-ph].
- [26] P. H. Chankowski and Z. Pluciennik, Phys. Lett. B **316**, 312 (1993) [arXiv:hep-ph/9306333]; K. S. Babu, C. N. Leung and J. T. Pantaleone, Phys. Lett. B **319**, 191 (1993) [arXiv:hep-ph/9309223]; S. Antusch, J. Kersten, M. Lindner and M. Ratz, Nucl. Phys. B **674**, 401 (2003) [arXiv:hep-ph/0305273].
- [27] S. Antusch, J. Kersten, M. Lindner, M. Ratz and M. A. Schmidt, JHEP **0503**, 024 (2005) [arXiv:hep-ph/0501272]; J. Chakraborty, A. Dighe, S. Goswami and S. Ray, arXiv:0812.2776 [hep-ph].
- [28] C. Hagedorn, J. Kersten and M. Lindner, Phys. Lett. B **597**, 63 (2004) [arXiv:hep-ph/0406103]; M. Honda, S. Kaneko and M. Tanimoto, JHEP **0309**, 028 (2003) [arXiv:hep-ph/0303227].
- [29] U. Sarkar and S. K. Singh, Nucl. Phys. B **771**, 28 (2007) [arXiv:hep-ph/0608030].

m_D	leptogenesis	wash-out
$\begin{pmatrix} 0 & 0 & a_3 \\ 0 & b_2 & 0 \\ 0 & c_2 & c_3 e^{i\gamma_3} \end{pmatrix}$	$\mathcal{I}_{23}^\tau = c_2^2 c_3^2 \sin 2\gamma_3$	$\tilde{m}_2^\tau = \frac{c_2^2}{M_2}$
$\begin{pmatrix} 0 & a_2 & 0 \\ 0 & 0 & b_3 \\ 0 & c_2 & c_3 e^{i\gamma_3} \end{pmatrix}$	$\mathcal{I}_{23}^\tau = c_2^2 c_3^2 \sin 2\gamma_3$	$\tilde{m}_2^\tau = \frac{c_2^2}{M_2}$
$\begin{pmatrix} a_1 & 0 & 0 \\ 0 & 0 & b_3 \\ c_1 & 0 & c_3 e^{i\gamma_3} \end{pmatrix}$	$\mathcal{I}_{13}^\tau = c_1^2 c_3^2 \sin 2\gamma_3$	$\tilde{m}_1^\tau = \frac{c_1^2}{M_1}$
$\begin{pmatrix} 0 & 0 & a_3 \\ b_1 & 0 & 0 \\ c_1 & 0 & c_3 e^{i\gamma_3} \end{pmatrix}$	$\mathcal{I}_{13}^\tau = c_1^2 c_3^2 \sin 2\gamma_3$	$\tilde{m}_1^\tau = \frac{c_1^2}{M_1}$
$\begin{pmatrix} 0 & a_2 & 0 \\ b_1 & 0 & 0 \\ c_1 & c_2 e^{i\gamma_2} & 0 \end{pmatrix}$	$\mathcal{I}_{12}^\tau = c_1^2 c_2^2 \sin 2\gamma_2$	$\tilde{m}_1^\tau = \frac{c_1^2}{M_1}$
$\begin{pmatrix} a_1 & 0 & 0 \\ 0 & b_2 & 0 \\ c_1 & c_2 e^{i\gamma_2} & 0 \end{pmatrix}$	$\mathcal{I}_{12}^\tau = c_1^2 c_2^2 \sin 2\gamma_2$	$\tilde{m}_1^\tau = \frac{c_1^2}{M_1}$

Table 2: The Dirac mass matrix, the non-zero expressions relevant for leptogenesis and the corresponding wash-out factors. All these cases give m_ν with one vanishing eigenvalue and $(m_\nu)_{e\mu} = 0$. The right-handed neutrino mass matrix is always diagonal.

m_D	Leptogenesis	washout
$\begin{pmatrix} 0 & 0 & a_3 \\ 0 & b_2 & b_3 e^{i\beta_3} \\ 0 & c_2 & 0 \end{pmatrix}$	$\mathcal{I}_{23}^\mu = b_2^2 b_3^2 \sin 2\beta_3$	$\tilde{m}_2^\mu = \frac{b_2^2}{M_2}$
$\begin{pmatrix} 0 & a_2 & 0 \\ 0 & b_2 & b_3 e^{i\beta_3} \\ 0 & 0 & c_3 \end{pmatrix}$	$\mathcal{I}_{23}^\mu = b_2^2 b_3^2 \sin 2\beta_3$	$\tilde{m}_2^\mu = \frac{b_2^2}{M_2}$
$\begin{pmatrix} 0 & 0 & a_3 \\ b_1 & 0 & b_3 e^{i\beta_3} \\ c_1 & 0 & 0 \end{pmatrix}$	$\mathcal{I}_{13}^\mu = b_1^2 b_3^2 \sin 2\beta_3$	$\tilde{m}_1^\mu = \frac{b_1^2}{M_1}$
$\begin{pmatrix} a_1 & 0 & 0 \\ b_1 & 0 & b_3 e^{i\beta_3} \\ 0 & 0 & c_3 \end{pmatrix}$	$\mathcal{I}_{13}^\mu = b_1^2 b_3^2 \sin 2\beta_3$	$\tilde{m}_1^\mu = \frac{b_1^2}{M_1}$
$\begin{pmatrix} 0 & a_2 & 0 \\ b_1 & b_2 e^{i\beta_2} & 0 \\ c_1 & 0 & 0 \end{pmatrix}$	$\mathcal{I}_{12}^\mu = b_1^2 b_2^2 \sin 2\beta_2$	$\tilde{m}_1^\mu = \frac{b_1^2}{M_1}$
$\begin{pmatrix} a_1 & 0 & 0 \\ b_1 & b_2 e^{i\beta_2} & 0 \\ 0 & c_2 & 0 \end{pmatrix}$	$\mathcal{I}_{12}^\mu = b_1^2 b_2^2 \sin 2\beta_2$	$\tilde{m}_1^\mu = \frac{b_1^2}{M_1}$

Table 3: The Dirac mass matrix, the non-zero expressions relevant for leptogenesis and the corresponding wash-out factors. All these cases give m_ν with one vanishing eigenvalue and $(m_\nu)_{e\tau} = 0$. The right-handed neutrino mass matrix is always diagonal.

m_D	Leptogenesis	washout	$\tan^2 \theta_{23}$
$\begin{pmatrix} 0 & a_2 & a_3 e^{i\alpha_3} \\ 0 & 0 & b_3 \\ 0 & 0 & c_3 \end{pmatrix}$	$\mathcal{I}_{23}^e = a_2^2 a_3^2 \sin 2\alpha_3$	$\tilde{m}_2^e = \frac{a_2^2}{M_2}$	$\frac{c_3^2}{b_3^2}$
$\begin{pmatrix} 0 & a_2 & a_3 e^{i\alpha_3} \\ 0 & b_2 & 0 \\ 0 & c_2 & 0 \end{pmatrix}$	$\mathcal{I}_{23}^e = a_2^2 a_3^2 \sin 2\alpha_3$	$\tilde{m}_2^e = \frac{a_2^2}{M_2}$	$\frac{c_3^2}{b_2^2}$
$\begin{pmatrix} a_1 & 0 & a_3 e^{i\alpha_3} \\ 0 & 0 & b_3 \\ 0 & 0 & c_3 \end{pmatrix}$	$\mathcal{I}_{13}^e = a_1^2 a_3^2 \sin 2\alpha_3$	$\tilde{m}_1^e = \frac{a_1^2}{M_1}$	$\frac{c_3^2}{b_3^2}$
$\begin{pmatrix} a_1 & 0 & a_3 e^{i\alpha_3} \\ b_1 & 0 & 0 \\ c_1 & 0 & 0 \end{pmatrix}$	$\mathcal{I}_{13}^e = a_1^2 a_3^2 \sin 2\alpha_3$	$\tilde{m}_1^e = \frac{a_1^2}{M_1}$	$\frac{c_1^2}{b_1^2}$
$\begin{pmatrix} a_1 & a_2 e^{i\alpha_2} & 0 \\ 0 & b_2 & 0 \\ 0 & c_2 & 0 \end{pmatrix}$	$\mathcal{I}_{12}^e = a_1^2 a_2^2 \sin 2\alpha_2$	$\tilde{m}_1^e = \frac{a_1^2}{M_1}$	$\frac{c_2^2}{b_2^2}$
$\begin{pmatrix} a_1 & a_2 e^{i\alpha_2} & 0 \\ b_1 & 0 & 0 \\ c_1 & 0 & 0 \end{pmatrix}$	$\mathcal{I}_{12}^e = a_1^2 a_2^2 \sin 2\alpha_2$	$\tilde{m}_1^e = \frac{a_1^2}{M_1}$	$\frac{c_1^2}{b_1^2}$

Table 4: The Dirac mass matrix, the non-zero expressions relevant for leptogenesis, the corresponding wash-out factors and the relevant part of the invariant for CP violation in neutrino oscillations. All these cases give m_ν with scaling property between 2nd and 3rd column, i.e., an inverted ordering with $m_3 = U_{e3} = 0$. The right-handed neutrino mass matrix is always diagonal.